

11 April 2019

A point in *rectangular* coordinates (x, y) can be written in *polar* coordinates (r, θ) , and vice versa. The correspondence is given by:

$$\begin{aligned} (x, y) &\rightarrow \left(\sqrt{x^2 + y^2}, \operatorname{atan2}(y, x) \right), \\ (r, \theta) &\rightarrow (r \cos(\theta), r \sin(\theta)), \end{aligned} \quad \operatorname{atan2}(y, x) = \begin{cases} \arctan(y/x) & \text{if } x > 0, \\ \arctan(y/x) + \pi & \text{if } x < 0, y \geq 0, \\ \arctan(y/x) - \pi & \text{if } x < 0, y < 0, \\ \pi/2 & \text{if } x = 0, y > 0, \\ -\pi/2 & \text{if } x = 0, y < 0, \\ 0 & \text{if } x = 0, y = 0. \end{cases}$$

1. **Warm up:** Convert the coordinates on the left to polar (r, θ) and the ones on the right to rectangular (x, y) .

(a) $(0, 0)$

(f) $(0, 0)$

(b) $(1, 0)$

(g) $(1, 0)$

(c) $(0, 1)$

(h) $(0, \pi)$

(d) $(-1, -1)$

(i) $(1, \pi)$


(e) $(-7, 24)$

(j) $(41\pi/4, 22\pi/3)$

2. Draw the given polar curves $r = f(\theta)$ on the graphs below.


(a)

$r = 3$




(c)

$r = 2/\cos(\theta)$




(b)

$r = \theta/\pi$




(d)

$r = 3 + \cos(3\theta)$



3. This question is about the folium of Descartes, the curve shown below. Its equation is $x^3 + y^3 = 3axy$, where $a \neq 0$ is a constant.



- (a) Show that for $t \neq -1, 0$, the line $y = tx$ intersects the folium at the origin and at one other point P . Express the coordinates of P in terms of t . Use this to obtain a parametrization of the folium almost everywhere.
- (b) Describe for which values of t the parametrization you found above describes the curve in quadrants I, II, and IV. Note $t = -1$ is a point of discontinuity of the parametrization.
- (c) Calculate dy/dx as a function of t and find the points with horizontal or vertical tangent.
- (d) Find a polar equation $r = f(\theta)$ of the folium.